

JOURNAL OF MATHEMATICAL ANALYSIS AND APPLICATIONS 33, 562–573 (1971)

## Existence of a Solution of a Boundary Value Problem for a Class of Discontinuous Nonlinear Systems

JAGDISH CHANDRA\*

*U. S. Army Research Office, Durham, N. C. 27706*

AND

B. A. FLEISHMAN†

*Department of Mathematics, Rensselaer Polytechnic Institute, Troy, New York**Submitted by R. Bellman*

### 1. INTRODUCTION

In this paper we investigate periodic solutions and related boundary value problems for a family of nonlinear differential equations in which the non-linearity may be discontinuous (and/or unbounded). Specifically, we establish the existence of harmonic solutions for equations of the type

$$\frac{d^2x}{dt^2} + x + f(x) = a \cos \omega t, \quad (DE, f)$$

where the function  $f$  is odd, satisfies  $xf(x) \geq 0$  for all  $x$  and has a jump discontinuity. All quantities are real-valued, and the constants  $a$  and  $\omega$  are assumed always to satisfy  $a < 0$  and  $\omega > 1$ .

What we ordinarily mean by a harmonic (periodic) solution of  $(DE, f)$  is a solution with period  $2\pi/\omega$  which is exactly out of phase (in phase when  $0 < \omega < 1$ ) with the forcing function  $a \cos \omega t$ . Now it is easily verified (see [5]) that a solution of the boundary value problem consisting of  $(DE, f)$  and the boundary conditions

$$\dot{x}(0) = 0, \quad x(\tau) = 0, \quad (1)$$

\* Research supported by the Department of Army Under Project No. IT061101 A91A.

† Research supported by Contract AF49(638)-1661 between the Office of Scientific Research of the U. S. Air Force and Rensselaer Polytechnic Institute.

(where  $\tau = \pi/2\omega$ ), when extended for all  $t$ , yields such a harmonic (periodic) solution of  $(DE, f)$ . Because it is the boundary value problem, denoted henceforth by  $BVP(f)$ , which is treated directly below, and because of the relationship between  $BVP(f)$  and the periodicity problem, *in this paper by a harmonic solution of  $(DE, f)$  we shall mean a solution of  $BVP(f)$  which is positive for  $0 \leq t < \tau$ .*

When  $f$  is uniformly bounded and smooth (except perhaps for a jump at  $x = 0$ ),  $(DE, f)$  is known [3] to possess a harmonic solution. The restriction that  $f$  experience a jump only at the origin is crucial in [3]; in this case a new (unsymmetric) smooth nonlinearity is constructed which coincides with  $f$  for  $x > 0$ . This technique does not work, however, when  $f$  has a jump discontinuity at points other than at  $x = 0$ . Our principal purpose here is to consider this problem. Further, it is shown that the restriction that  $f$  be uniformly bounded may be relaxed considerably: in Section 4, we consider a class of unbounded nonlinearities which satisfy suitable growth conditions.

The existence of a solution of  $BVP(f)$  is established below by use of the well-known point-mapping method; to employ this method we need to know that, for each value of the real parameter  $\alpha$  in a certain interval, the initial value problem  $(DE, f)$ ,  $x(0) = \alpha$ ,  $\dot{x}(0) = 0$  possesses a unique solution on  $[0, \tau]$  which depends continuously on  $\alpha$ . This initial value problem is treated in Section 5.

For ease of exposition we have restricted ourselves to functions  $f$  with a single discontinuity. Similar results can be obtained for a broader class of problems, involving nonlinearities with multiple discontinuities. Such problems will be discussed elsewhere. A particular case of multiple discontinuities, in which the nonlinearity is piecewise constant, is treated in [5] by a special technique.

## 2. PRELIMINARIES

Let  $b = a/(1 - \omega^2)$ .

$F$  will denote the class of functions  $f$  with the following properties:  $f$  is odd,  $f(x) \geq 0$  when  $x > 0$ ,  $f(x) \neq 0$ , and  $f$  is continuous for  $x \geq 0$  except perhaps at  $x = \sigma$  ( $0 < \sigma < b$ ), where it has a jump discontinuity.

In Sections 3, 4 and 5 we introduce subsets of  $F$  denoted, respectively, by  $F_M$  ( $f$  uniformly bounded by a positive constant  $M$ ),  $F_g$  ( $f$  unbounded but subject to a growth restriction), and  $F_L$  ( $f \in F_M \cup F_g$  but in addition satisfying a Lipschitz condition in each interval of continuity).

When  $f$  is piecewise continuous we mean by a solution of  $(DE, f)$  on an interval  $I$  a function  $x(t)$  which is continuously differentiable on this interval, possesses a piecewise continuous second derivative and satisfies  $(DE, f)$

wherever the second derivative exists. In case  $f$  is continuous a solution is to have a continuous second derivative and satisfy  $(DE, f)$  at all points of  $I$ .

Let  $IVP(\alpha, f)$  denote the initial value problem of solving  $(DE, f)$  subject to the initial conditions

$$x(0) = \alpha, \quad \dot{x}(0) = 0. \quad (2)$$

By use of the method of variation of parameters, a solution of  $IVP(\alpha, f)$  on an interval  $I$  containing the origin is easily seen to satisfy

$$x(t) = (\alpha - b) \cos t + b \cos \omega t - \int_0^t \sin(t-s)f(x(s)) ds, \quad (t \in I). \quad (3)$$

Conversely, a continuous solution of (3) on  $I$  is a solution of  $IVP(\alpha, f)$  on  $I$ .

Let  $x(t; \alpha, f)$  denote a solution of  $IVP(\alpha, f)$  which is known to exist and be unique on  $[0, \tau]$ . (When  $f$  is Lipschitzian, such a unique solution is known to exist (for instance, see [4]); this is less evident, however, when  $f$  is piecewise continuous. We establish in Section 5 the existence, uniqueness and continuous dependence on initial data of solutions of  $IVP(\alpha, f)$  when  $f$  belongs to a subset of our class  $F$  of piecewise continuous nonlinearities.)

Let  $A$  be a set of values of  $\alpha$  for which (unique)  $x(t; \alpha, f)$  exists. We define on  $A$  a mapping  $T$  by the formula

$$T(\alpha) = x(\tau; \alpha, f).$$

Clearly  $x(t; \alpha, f)$  will be a harmonic solution of  $(DE, f)$  for those values of  $\alpha$  for which  $T(\alpha) = 0$  and  $x(t; \alpha, f) > 0$  for  $0 \leq t < \tau$ . Any such value of  $\alpha$  will henceforth be called a *harmonic point*.

The following theorem is a restatement, under weaker assumptions, of one appearing originally in [2] and [3].

**COMPARISON THEOREM.** *Suppose that  $f_1$  and  $f_2$  are piecewise continuous functions and that  $f_1(r) \geq f_2(s)$  for all  $r > 0, s > 0$ . Let  $\alpha_1$  be a harmonic point of  $(DE, f_1)$  and  $\alpha_2$  a harmonic point of  $(DE, f_2)$ . Then*

$$\alpha_1 \geq \alpha_2.$$

*Proof.* Let  $x_i(t; a_i, f_i)$  be a harmonic solution of  $(DE, f_i)$  ( $i = 1, 2$ ). Then, using (3), the harmonic points  $\alpha_i$  ( $i = 1, 2$ ) are given by

$$\alpha_i = b + \sec \tau \int_0^\tau \sin(\tau - s) f_i(x_i(s)) ds,$$

so that

$$\alpha_1 - \alpha_2 = \sec \tau \int_0^\tau \sin(\tau - s) \{f_1(x_1) - f_2(x_2)\} ds \geq 0.$$

Here we have used the facts that  $0 \leq \tau - s \leq \pi/2$  and  $x_i(s; \alpha_i, f_i) > 0$  ( $i = 1, 2$ ) for  $s \in (0, \tau)$ .

In the linear case,  $f(x) \equiv 0$ , the equation  $(DE, 0)$  has a unique harmonic point  $b = a/(1 - \omega^2)$ . The ideal relay equation

$$\frac{d^2x}{dt^2} + x + M \operatorname{sgn} x = a \cos \omega t, \quad (DE, M \operatorname{sgn})$$

where  $M$  is a positive constant and  $\operatorname{sgn} x = \text{signum } x = x/|x|$  ( $x \neq 0$ ), is known [7] to have a harmonic point, which we denote by  $d_M$ :

$$d_M = b + M(\sec \tau - 1). \quad (4)$$

According to the Comparison Theorem,  $d_M$  is the unique harmonic point of  $(DE, M \operatorname{sgn})$ .

### 3. EXISTENCE OF A HARMONIC SOLUTION: $f$ UNIFORMLY BOUNDED

In this section we assume always that  $f$  is contained in  $F_M$ , the subset of  $F$  consisting of functions which satisfy  $|f(x)| \leq M$  for all  $x$ , where  $M$  is a given positive constant.

When  $f \in F_M$  it is not hard to see (since  $f(x) \not\equiv 0$ ) that there is a subinterval of  $(0, d_M)$ , on which  $f(x) > 0$ , and there is also a subinterval of  $(0, d_M)$  on which  $f(x) < M$ . These facts are needed in Lemma 2 and Corollary 1 of Lemma 3.

From the Comparison Theorem follows immediately

LEMMA 1. *If  $\alpha$  is a harmonic point of  $(DE, f)$ ,*

$$b \leq \alpha \leq d_M.$$

Thus, for  $f \in F_M$  we know that we must look for a harmonic point (if any exists) in the interval  $[b, d_M]$ . To show that one does exist (that is, that there exists at least one harmonic solution) we use the well-known point-mapping method [1]; namely, we demonstrate that  $T$  is a continuous mapping and carries the interval  $[b, d_M]$  into an interval which contains zero in its interior (so that at least one point in  $[b, d_M]$  is mapped into zero under  $T$ ).

To this end we show below that  $T(d_M) > 0$ ,  $T(b) < 0$  and  $\dot{x}(t; \alpha, f) < 0$  on  $0 < t \leq \tau$  for any  $\alpha \in [b, d_M]$ . The continuity of  $T(\alpha)$  in  $\alpha$  follows from the continuous dependence of  $x(t; \alpha, f)$  on the initial datum  $\alpha$ , which is established in Section 5. These facts yield the desired result, Theorem 1.

In Lemma 2 and Lemma 3 we assume that for each  $\alpha \in [b, d_M]$  there exists a solution of  $IVP(\alpha, f)$  which is defined for all  $t \in [0, \tau]$ . In Section 5 we demonstrate that this is not an empty assumption.

LEMMA 2.  $T(d_M) = x(\tau; d_M, f) > 0$ .

*Proof.* Since  $|f(x)| \leq M$ ,

$$\begin{aligned} \int_0^\tau \sin(\tau - s) f(x(s)) ds &\leq M \int_0^\tau \sin(\tau - s) ds \\ &= M(1 - \cos \tau). \end{aligned}$$

But

$$\begin{aligned} x(\tau; d_M, f) &= (d_M - b) \cos \tau - \int_0^\tau \sin(\tau - s) f(x(s)) ds \\ &\geq (d_M - b) \cos \tau - M(1 - \cos \tau) \\ &= [d_M - b - M(\sec \tau - 1)] \cos \tau \\ &= 0. \end{aligned}$$

Thus  $x(\tau; d_M, f) \geq 0$ . Now suppose  $x(\tau; d_M, f) = 0$ . Then  $x(t; d_M, f)$  takes all values in  $0 \leq x \leq d_M$ , on a subinterval of which  $f(x) < M$ . Therefore, retracing the steps above,

$$\int_0^\tau \sin(\tau - s) f(x(s)) ds < M(1 - \cos \tau)$$

and

$$x(\tau; d_M, f) > 0.$$

COROLLARY. If  $\alpha$  is a harmonic point of  $(DE, f)$  then  $\alpha < d_M$ .

*Proof.* From Lemma 1,  $\alpha \leq d_M$ . But if  $\alpha = d_M$ ,  $x(\tau; d_M, f) = 0$ , contradicting Lemma 2.

LEMMA 3. Suppose  $b\omega \geq M$ . Then for  $\alpha \in [b, d_M]$ ,  $\dot{x}(t; \alpha, f) < 0$  on  $0 < t \leq \tau$ .

*Proof.*

$$\begin{aligned} \dot{x}(t; \alpha, f) &= (b - \alpha) \sin t - b\omega \sin \omega t - \int_0^t \cos(t - s) f(x(s)) ds \\ &\leq (b - \alpha) \sin t - b\omega \sin \omega t + M \int_0^t \cos(t - s) ds \\ &\leq (b - \alpha) \sin t - b\omega(\sin \omega t - \sin t) \\ &\leq -\eta(t) < 0 \end{aligned}$$

where

$$\eta(t) = b\omega(\sin \omega t - \sin t).$$

COROLLARY 1.  $T(b) = x(\tau; b, f) < 0$ .

*Proof.* On the contrary, assume that  $x(\tau; b, f) \geq 0$ . Then, since  $\dot{x}(t; b, f) < 0$ ,  $x(t; b, f) > 0$  for  $0 \leq t < \tau$ . This implies  $f(x(t)) > 0$ , so that from (3)

$$x(\tau; b, f) = - \int_0^\tau \sin(\tau - s) f(x(s)) ds < 0.$$

This contradiction completes the proof.

COROLLARY 2. If  $\alpha$  is a harmonic point of  $(DE, f)$ , then  $\alpha > b$ .

On the basis of the preceding lemmas, which establish the fact that the mapping  $T$  carries  $[b, d_M]$  into an interval containing zero in its interior, and in view of continuity of the solution  $x(t; \alpha, f)$  in  $\alpha$  (which will be demonstrated in Section 5), we have proved

THEOREM 1. Suppose  $f \in F_M$ . Then there exists an  $\alpha$ ,  $\alpha \in (b, d_M)$ , which is a harmonic point of  $(DE, f)$ , provided

$$b\omega \geq M.$$

#### 4. EXISTENCE OF A HARMONIC SOLUTION: $f$ UNBOUNDED

Now we extend the results of the preceding section to a class of unbounded nonlinearities. In this section it is assumed always that  $f$  belongs to  $F_g$ , the subset of  $F$  consisting of functions  $f$  which obey the following growth condition: there exists a positive number  $\bar{x}$  (where  $\bar{x} > b > \sigma > 0$ ) such that for  $x > 0$

$$\begin{aligned} f(x) &\leq f(\bar{x}) = \frac{\bar{x} - b}{\sec \tau - 1}, & (x \leq \bar{x}), \\ f(x) &< \frac{x - b}{\sec \tau - 1}, & (x > \bar{x}). \end{aligned} \tag{5}$$

To motivate this extension, let us consider an arbitrary  $f \in F$ . For a given  $\alpha \geq b$ , suppose  $x(t; \alpha, f)$  is a harmonic solution of  $(DE, f)$ . Since  $f(x(t)) \geq 0$  for  $0 \leq t \leq \tau$ , it follows from  $\dot{x}(t; \alpha, f) < 0$  on  $(0, \tau]$  (modifying slightly the proof of Lemma 3) that  $x(t; \alpha, f) < \alpha$  on  $(0, \tau)$ . Define the odd truncated function  $f_0$  by

$$f_0(x) = \begin{cases} f(x), & 0 \leq x \leq \alpha \\ f(\alpha), & x > \alpha. \end{cases}$$

Since  $x(t; \alpha, f)$  takes values only in  $-\alpha \leq x \leq \alpha$ , where  $f$  and  $f_0$  are identical,  $x(t; \alpha, f)$  is a harmonic solution of

$$\frac{d^2x}{dt^2} + x + f_0(x) = a \cos \omega t \quad (DE, f_0)$$

as well as of  $(DE, f)$ . Since  $|f_0(x)| \leq f(\alpha)$ , we find, from the Comparison Theorem of Section 2, using  $M = f(\alpha)$ , that

$$b \leq \alpha \leq b + f(\alpha) [\sec \tau - 1]. \quad (6)$$

Inequality (6) imposes a necessary condition to be satisfied by the amplitude  $\alpha$  of a harmonic solution (if any exists) of  $(DE, f)$ . Thus, if for all  $x$  greater than some  $\bar{x} > 0$ ,  $x > b + f(x) [\sec \tau - 1]$ , there cannot exist a harmonic solution with amplitude greater than  $\bar{x}$ . This motivates the definition of the set  $F_g$ .

On the basis of the preceding discussion, we have

LEMMA 6. Suppose  $f \in F_g$ . If  $\alpha$  is a harmonic point of  $(DE, f)$ , then

$$b \leq \alpha \leq \bar{x}. \quad (7)$$

Thus, in order to investigate harmonic solutions of  $(DE, f)$ , for  $f \in F_g$ , it is sufficient to consider solutions of  $IVP(\alpha, f)$  for  $b \leq \alpha \leq \bar{x}$ .

Define the truncated function  $\bar{f}$  by

$$\bar{f}(x) = \begin{cases} f(x), & 0 \leq x \leq \bar{x} \\ \frac{\bar{x} - b}{\sec \tau - 1}, & x > \bar{x}. \end{cases} \quad (8)$$

Clearly  $|\bar{f}(x)| \leq (\bar{x} - b)/(\sec \tau - 1)$ , and Lemmas 2 and 3 can be restated in terms of  $\bar{f}$ , and thus in terms of  $f$ . We have, for instance,

LEMMA 2'. Let  $x(t; \bar{x}, f)$  be a solution of  $IVP(\bar{x}; f)$ . Then  $x(\tau; \bar{x}, f) > 0$ . If  $\alpha$  is a harmonic point of  $(DE, f)$ , then  $\alpha < \bar{x}$ .

We shall merely state the existence result.

THEOREM 2. Let  $f \in F_g$ . Then there exists an  $\alpha \in (b, \bar{x})$  which is a harmonic point of  $(DE, f)$  provided

$$b\omega \geq \frac{\bar{x} - b}{\sec \tau - 1}.$$

### 5. EXISTENCE, UNIQUENESS AND CONTINUOUS DEPENDENCE OF A SOLUTION OF $IVP(\alpha, f)$

Let  $F_L$  be the subset of  $F$  consisting of those elements in  $F_M$  or  $F_g$  which, except at  $x = \pm \sigma$ , are Lipschitzian: for  $x \geq 0$

$$f(x) = \begin{cases} f_1(x), & x \leq \sigma \\ f_2(x), & x \geq \sigma \end{cases}$$

where  $f_1$  and  $f_2$  are Lipschitz continuous on their respective domains of definition. Obviously  $F_L \subset (F_M \cup F_g)$ .

Throughout this section it is assumed that  $\alpha \in [b, \bar{x}]$ ,  $\bar{x} > b > \sigma > 0$ , and that  $f \in F_L$ . Calculations and results are stated explicitly for members  $f$  of  $F_g$  (which satisfy the growth restriction (5)). They are easily modified for members  $f$  of  $F_M$  (uniformly bounded).

We work with the trajectories of solutions in  $t, x, \dot{x}$ -space (phase space). Given the constant  $N > 0$  (which for the moment is arbitrary), let

$$D = D_1 \cup D_2 \cup D_3$$

where

$$D_1 = \{(t, x, \dot{x}) \mid 0 \leq t \leq \tau, \sigma \leq x \leq \bar{x}, |\dot{x}| \leq N\}$$

$$D_2 = \{(t, x, \dot{x}) \mid 0 \leq t \leq \tau, -\sigma \leq x \leq \sigma, |\dot{x}| \leq N\}$$

$$D_3 = \{(t, x, \dot{x}) \mid 0 \leq t \leq \tau, -\bar{x} \leq x \leq -\sigma, |\dot{x}| \leq N\}.$$

We shall prove that for each  $\alpha \in [b, \bar{x}]$  the solution of  $IVP(\alpha, f)$  can be continued uniquely up to  $t = \tau$  and that this solution depends continuously on the initial datum  $\alpha$ .

Since in  $D_1$   $f$  is Lipschitz continuous, there exists a  $t_1$  ( $0 < t_1 \leq \tau$ ) such that for  $0 < t < t_1$ ,  $IVP(\alpha, f)$  has a unique solution whose trajectory is in  $D_1$ . This trajectory approaches the boundary of  $D_1$  as  $t \rightarrow t_1$  (for example, see [8]). Let us denote this solution by  $x(t; \alpha, f)$ . Then  $x(t; \alpha, f)$  satisfies the integral equation (3) (with  $f$  replaced by  $f_2$ ) on  $[0, t_1]$ . For  $0 < t < t_1$

$$\begin{aligned} \dot{x}(t; \alpha, f) &= -(\alpha - b) \sin t - b\omega \sin \omega t - \int_0^t \cos(t-s) f_2(x(s)) ds \\ &\leq -(\alpha - b) \sin t - b\omega \sin \omega t + \frac{\bar{x} - b}{\sec \tau - 1} \sin t \\ &\leq -\eta(t) < 0, \end{aligned}$$

where  $\eta(t) = b\omega(\sin \omega t - \sin t)$ , provided

$$b\omega \geq \frac{\bar{x} - b}{\sec \tau - 1}. \quad (9)$$



(For the remainder of this section we assume that (9) holds.) Further,

$$\begin{aligned} -\dot{x}(t; \alpha, f) &\leq (\alpha - b) \sin t + b\omega \sin \omega t + \frac{\bar{x} - b}{\sec \tau - 1} \int_0^t \cos(t-s) ds \\ &\leq (\bar{x} - b) \sin \tau + b\omega + \frac{\bar{x} - b}{1 - \cos \tau} (\sin \tau \cos \tau) \\ &= b\omega + \frac{\bar{x} - b}{1 - \cos \tau} (\sin \tau). \end{aligned}$$

Therefore, if we choose

$$N > b\omega + \frac{\bar{x} - b}{1 - \cos \tau} (\sin \tau),$$

$$0 > \dot{x}(t; \alpha, f) > -N.$$

Thus, the trajectory of solution  $x(t; \alpha, f)$ , for  $0 \leq t < t_1$ , is bounded away from the boundaries  $\dot{x} = \pm N$  and remains below the plane  $x = \bar{x}$ . Hence, as  $t \rightarrow t_1$  the trajectory either approaches the boundary  $t = \tau$  (in which case  $t_1 = \tau$ ) or the boundary  $x = \sigma$ . Now if  $t_1 = \tau$  we have a well defined solution of  $(DE, f)$  on  $0 \leq t \leq \tau$  which depends continuously on  $\alpha$ . When  $t_1 < \tau$  define

$$v_1(t_1, \alpha) = \lim_{t \rightarrow t_1^-} \dot{x}(t; \alpha, f) < 0.$$

Thus  $x(t; \alpha, f)$  is not tangent to the plane  $x = \sigma$  at  $t = t_1$ . Clearly  $v_1$  is a continuous function of  $(t_1, \alpha)$ . Now  $t_1$  is obtained by solving the functional equation

$$x(t_1; \alpha, f) = \sigma.$$

Since  $x(t; \alpha, f)$  is a strictly decreasing function of  $t$ , there exists a unique solution  $t_1 = u_1(\alpha)$ , where  $u_1(\alpha)$  is a continuous function of  $\alpha$ .

We now continue this solution for  $t > t_1$  as the unique solution of the initial value problem

$$\frac{d^2x}{dt^2} + x + f_1(x) = a \cos \omega t$$

$$x(t_1) = \sigma, \quad \dot{x}(t_1) = v_1 < 0$$

in  $D_2$ . Since  $f_1$  is Lipschitz continuous in  $D_2$ , there exists a  $t_2$  ( $t_1 < t_2 \leq \tau$ ) such that the unique continuation remains in  $D_2$  for  $t_1 < t < t_2 \leq \tau$  and depends continuously on the "initial" data  $t_1, v_1$ . Since these "initial" data themselves are continuous functions of  $\alpha$ , we are assured of continuous dependence on  $\alpha$  of the extended solution on  $0 \leq t < t_2 \leq \tau$ .

In order to gain insight into subsequent continuation of this solution, let us consider the comparison problem

$$\frac{d^2x}{dt^2} + x + \frac{\bar{x} - b}{\sec \tau - 1} = a \cos \omega t,$$

$x(0) = b$ ,  $\dot{x}(0) = 0$ . The unique solution of this problem is given by

$$x_\ell(t) = \frac{\bar{x} - b}{\sec \tau - 1} (\cos t - 1) + b \cos \omega t.$$

On their common interval of existence,  $x_\ell(t)$  provides a lower bound for  $x(t; \alpha, f)$ :

$$\begin{aligned} x(t; \alpha, f) - x_\ell(t) &= (\alpha - b) \cos t - \frac{\bar{x} - b}{\sec \tau - 1} (\cos t - 1) \\ &\quad - \int_0^t \sin(t-s) f(x(s)) ds \\ &\geq (\alpha - b) \cos t - \frac{\bar{x} - b}{\sec \tau - 1} (\cos t - 1) \\ &\quad + \frac{\bar{x} - b}{\sec \tau - 1} (\cos t - 1) \\ &= (\alpha - b) \cos t \\ &\geq 0. \end{aligned}$$

Since  $x_\ell(t)$  is a strictly decreasing function of  $t$  on  $[0, \tau]$ ,  $x(t; \alpha, f)$  remains above the boundary  $x = -\sigma$  for  $0 \leq t \leq \tau$  (and therefore  $t_2 = \tau$ ) if

$$x_\ell(\tau) = \frac{\bar{x} - b}{\sec \tau - 1} (\cos \tau - 1) = -(\bar{x} - b) \cos \tau > -\sigma$$

or

$$\sigma > (\bar{x} - b) \cos \tau. \quad (10)$$

*Remark.* If we assume that

$$b > \sigma > b\omega(1 - \cos \tau)$$

then (9) implies (10); for from (9)

$$\bar{x} - b \leq b\omega(\sec \tau - 1) < \sigma \sec \tau.$$

If (10) does not hold, the trajectory for  $x(t; \alpha, f)$  may approach the boundary  $x = -\sigma$  as  $t \rightarrow t_2$ . It is easily verified that

$$v_2(t_2, \alpha) = \lim_{t \rightarrow t_2-0} \dot{x}(t; \alpha, f) < 0.$$

We now continue the solution for  $t > t_2$  as the unique solution of the initial value problem

$$\frac{d^2x}{dt^2} + x + f_2(x) = a \cos \omega t$$

satisfying  $x(t_2) = -\sigma$ ,  $\dot{x}(t_2) = v_2 < 0$ . The trajectory thus obtained remains above the plane  $x = -\bar{x}$  (and therefore reaches the plane  $t = \tau$ ); for, from the comparison problem,

$$x(t; \alpha, f) \geq x_c(\tau) = -(\bar{x} - b) \cos \tau > -\bar{x}.$$

The above discussion is summarized in

**THEOREM 3.** *Suppose  $f \in F_L \cap F_g$ , where  $0 < \sigma < b < \bar{x}$ . Then for each  $\alpha \in [b, \bar{x}]$  there exists a unique solution  $x(t; \alpha, f)$  of the initial value problem  $IVP(\alpha, f)$  on  $0 \leq t \leq \tau$  which depends continuously on  $\alpha$ , provided*

$$b\omega \geq \frac{\bar{x} - b}{\sec \tau - 1}.$$

*In phase space the trajectory of this solution crosses each plane of discontinuity  $x = \pm \sigma$  at most once.*

If further  $\sigma > b\omega(1 - \cos \tau)$  the unique solution of  $IVP(\alpha, f)$  remains above the plane  $x = -\sigma$  for  $0 \leq t \leq \tau$ ; that is, the trajectory intersects at most one plane of discontinuity of  $f$ . (If  $f \in F_L \cap F_M$ , replace  $\bar{x}$  above by  $d_M$ .)

## REFERENCES

1. A. A. ANDRONOV, A. A. VITT, AND S. E. KHAIKIN, "Theory of Oscillators," Pergamon Press, New York, 1966. (Translated from Russian.)
2. O. BENEDIKTSSON, Forced oscillations in second order systems with bounded nonlinearities, Doctoral Dissertation, Rensselaer Polytechnic Institute, 1965.
3. O. BENEDIKTSSON AND B. A. FLEISHMAN, Forced oscillations in second order systems with bounded nonlinearities, in "Proc. Internat. Symposium on Differential Equations and Dynamical Systems," pp. 497-503, Academic Press, New York, 1967.
4. L. CESARI, "Asymptotic Behavior and Stability Problems in Ordinary Differential Equations," Springer-Verlag, Berlin, 1963.
5. J. CHANDRA AND B. A. FLEISHMAN, Positivity and comparison results for nonlinear boundary value problems and related periodic solutions, *J. Math. Anal. Appl.* **24** (1968), 545-554.
6. J. CHANDRA AND B. A. FLEISHMAN, A canonical representation for solutions of a class of nonlinear problems, *Internat. J. Nonlinear Mech.* **5** (1970), 63-79.

7. B. A. FLEISHMAN, Harmonic and subharmonic responses of an on-off control system to sinusoidal inputs, *J. Franklin Inst.* **270** (1960), 99–113.
8. W. HUREWICZ, "Lectures on Ordinary Differential Equations," Technology Press of M.I.T., also John Wiley and Sons, London, 1958.